EPI SCIENTIFIC REFERENCES AVAILABLE

- 1. Amiali, M. et al. (2007). Synergistic effect of temperature and pulsed electric field on inactivation of *Escherichia coli* O157:H7 and *Salmonella enteritidis* in liquid egg yolk. Journal of Food Engineering <u>79</u>: 689-694
- Banerjee, P. et al. (2012). Maximizing carbon dioxide content of shell eggs by rapid cooling treatment and its effect on shell egg quality. Poultry Science <u>91:</u> 1444-1453
- Bengoechea, C. et al. (2009). Influence of semicontinuous processing on the rheology and droplet size distribution of mayonnaise-like emulsion. Food Science & Technology International <u>15:</u> 367-373
- Buckman, J. et al. (2002). Note. The effect of a commercial extended egg albumen on the microstructure of Icing. Food Science & Technology International <u>8</u>: 109-115
- 5. Campbell, L. et al. (2005). Heat stability and emulsifying ability of whole egg and egg yolk as related to heat treatment. Food Hydrocolloids <u>19</u>: 533-539
- 6. Coimbra, J.S.R. et al. (2006). Density, heat capacity and thermal conductivity of liquid egg products. Journal of Food Engineering <u>74:</u> 186-190
- 7. Daimer, K. et al. (2008). Oil-in-water emulsion properties of egg yolk: Effect of enzymatic modification by phospholipase A₂. Food Hydrocolloids <u>23:</u> 1366-1373
- De Souza, P.M. et al. (2011). Effects of UV-C on physiochemical quality attributes and *Salmonella enteritidis* inactivation in liquid egg products. Food Control <u>22:</u> 1385-1392
- De Souza, P.M. et al. (2012). Rheological properties and protein quality of UV-C processed liquid egg products. Food Hydrocolloids doi:10.1016/j.foodhyd.2012.05.013
- Faeste, C.K. et al. (2007). Extractability, stability, and allergenicity of egg white proteins in differently heat-processed foods. Journal of the AOAC International <u>90:</u> 427-436
- Ganasen, P. et al. (2011). Effects of three cations on the stability and microstructure of protein aggregate from duck egg white under alkaline condition. Food Science & technology International <u>17:</u> 343-349

- 12. Garcia-Gonzalez, L. et al. (2009). Inactivation of naturally occurring microorganisms in liquid whole egg using high pressure carbon dioxide processing as an alternative to heat pasteurization. Journal of Supercritical Fluids <u>51:</u> 74-82
- 13. Geera, B. et al. (2011). A comprehensive evaluation of egg and egg replacers on the product quality of muffins. Journal of Food Quality <u>34:</u> 333-342
- 14. Grijspeerdt, K. et al. (2003). Inactivation of *Salmonella enteritidis* during boiling of eggs. International Journal of Food Microbiology <u>82:</u> 13-24
- 15. Guerrero, A. et al. (2000). Linear and non-linear viscoelasticity of low-incholesterol mayonnaise. Food Science & Technology International <u>6:</u> 165-172
- Guilmineau, F. et al. (2006). Impact of a thermal treatment on the emulsifying properties of egg yolk. Part 1: Effect of the heating time. Food Hydrocolloids <u>20:</u> 1105-1113
- Guilmineau, F. et al. (2006). Impact of a thermal treatment on the emulsifying properties of egg yolk. Part 2: Effect of the environmental conditions. Food Hydrocolloids <u>20:</u> 1114-1123
- 18. Guilmineau, F. et al. (2007). Influence of a thermal treatment on the functionality of hen's egg yolk in mayonnaise. Journal of Food Engineering <u>78:</u> 648-654
- Gujral, H.S. et al. (2003). Effect of sodium lauryl sulphate on the texture of sponge cake. Food Science & Technology International <u>9:</u> 89-93
- Huang, E., et al. (2006). Inactivation of Salmonella enteritidis in liquid whole egg using combination treatments of pulsed electric filed, high pressure and ultrasound. Biosystems Engineering <u>94:</u> 403-413
- 21. Huang, Q. et al. (2012). Estimation of egg freshness using S-ovalbumin as an indicator. Poultry Science <u>91:</u> 739-743
- Ibarz, A. et al. (1999). Texture of egg gels induced by high hydrostatic pressure (In Spanish). Food Science & Technology International <u>5:</u> 191-201
- 23. Jaekel, T. et al. (2008). Preserving functional properties of hen's egg yolk during freeze-drying. Journal of Food Engineering <u>87:</u> 522-526
- 24. Jones, D.R. et al. (2005). A survey of common practices in shell egg processing facilities and water use. International Journal of Poultry Science <u>4</u>: 734-736

- 25. Jones, D.R et al. (2007). Pathogen prevalence and microbial levels associated with restricted shell eggs. Journal of Food Protection <u>70:</u> 2004-2007
- Jordan, J.S. et al. (2011). A mathematical model of inactivation kinetics for a fourstrain composite of Salmonella enteritidis and Oranienburg in commercial liquid egg yolk. Food Microbiology <u>28:</u> 67-75
- Kaewmanee, T. et al. (2009). Changes in chemical composition, physical properties and microstructure of duck egg as influenced by salting. Food Chemistry <u>112</u>: 560-569
- 28. Kaewmanee, T. et al. (2011). Effect of NaCl on thermal aggregation of egg white proteins from duck egg. Food Chemistry <u>125</u>: 706-712
- Kassis, N. et al. (2010). Development of nutraceutical egg products with omega-3rich oils. LWT – Food Science and Technology <u>43:</u> 777-783
- Laca, A. et al. (2010). A method of egg yolk fractionation. Characterization of fractions. Food Hydrocolloids <u>24:</u> 434-443
- Lechevalier, V. et al. (2007). Egg white drying: Influence of industrial processing steps on protein structure and functionalities. Journal of Food Engineering <u>83:</u> 404-413
- 32. Maňas, P. et al. (2003). Survival of *Salmonella senftenberg* 775W to current liquid whole egg pasteurization treatments. Food Microbiology <u>20:</u> 593-600
- McQuestin, O.J. et al. (2010). Kinetics of growth and inactivation of Salmonella enterica serotype typhimurium DT104 in pasteurised liquid egg products. Food Microbiology <u>27:</u> 396-402
- Monfort, S. et al. (2010). Inactivation of Salmonella typhimurium and Staphylococcus aureus by pulsed electric fields in liquid whole egg. Innovative Food Science and Emerging Technologies <u>11:</u> 306-313
- Monfort, S. et al. (2011). Design of a combined process for the inactivation of Salmonella enteritidis in liquid whole egg at 55° C. International Journal of Food Microbiology <u>145</u>: 476-482
- 36. Monfort, S. et al. (2012). Inactivation of *Salmonella spp.* in liquid whole egg using pulsed electric fields, heat, and additives. Food Microbiology <u>30:</u> 393-399

EPI Scientific literature available - Listed by alphabetic order – 15-August-2012

- Musgrove, M.T. et al. (2012). Evaluation of an alcohol-based sanitozer spray's bactericidal effects on *Salmonella* inoculated onto stainless steel and shell egg processing equipment. International Journal of Poultry Science <u>11:</u> 92-95
- Nemeth, C.S. et al. (2011). Thermal destruction of *Listeria monocytogenes* in liquid egg products with heat treatment at lower temperature and longer than pasteurization. African Journal of Food Science <u>5</u>: 161-167
- Riscardo, M.A. et al. (2003). Influence of composition of emulsifier blends on the rheological properties of salad dressing-type emulsions. Food Science & Technology International <u>9:</u> 53-63
- 40. Rocculi, P. et al. (2011). Modified atmosphere packaging of hen table eggs: Effects on functional properties of albumen. Poultry Science <u>90:</u> 1791-1798
- Rossi, M. et al. (2010). Functional properties of pasteurised liquid whole egg products as affected by the hygienic quality of raw eggs. LWT – Food Science and Technology <u>43:</u> 436-441
- 42. Sampedro, F. et al. (2006). Review: Application of pulsed electric fields in egg and egg derivatives. Food Science & Technology International <u>12:</u> 397-406
- 43. Silversides, F.G. et al. (2004). The relationship among measures of egg albumen height, pH, and whipping volume. Poultry Science <u>83:</u> 1619-1623
- Talansier, E. et al. (2009). Optimization of dry heat treatment of egg white in relation to foam and interfacial properties. LWT – Food Science and Technology <u>42:</u> 496-503
- 45. Tedeso, J. et al. 92006). Fatty acid profiles of processed chicken egg yolks. Journal of Agricultural and Food Chemistry <u>54</u>: 6255-6260
- 46. Unluturk, S. et al. (2008). Use of UV-C radiation as a non-thermal process for liquid egg products (LEP). Journal of Food Engineering <u>85:</u> 561-568
- Unluturk, S. et al. (2010). Modelling inactivation kinetics of liquid egg white exposed to UV-C irradiation. International Journal of Food Microbiology <u>142:</u> 341-347
- Van der Plancken, I. et al. (2007). Foaming properties of egg white proteins affected by heat or high pressure treatment. Journal of Food Engineering <u>78:</u> 1410-1426

EPI Scientific literature available - Listed by alphabetic order – 15-August-2012